



ELSEVIER

Discrete Mathematics 236 (2001) 95–109

DISCRETE
MATHEMATICSwww.elsevier.com/locate/disc

New classes of perfectly orderable graphs

J.-L. Fouquet^a, F. Roussel^b, P. Rubio^b, H. Thuillier^{b,*}^a*IUT Mesures Physiques, Université du Maine, Avenue O. Messiaen 72017 Le Mans Cedex, France*^b*Université d'Orléans, L.I.F.O., Bat IIIA, Rue Leopard de Vinci, B.P. 6759,
F-45067 Orléans Cedex 2, France*

Received 17 November 1997; revised 28 July 1999; accepted 5 March 2000

Abstract

This paper generalizes previous works on perfectly orderable graphs by Olariu (Discrete Math. 113 (1992) 143) and by Hoàng et al. (Discrete Math. 102 (1992) 67). Chvátal defined a graph to be perfectly orderable (V. Chvátal, in: C. Berge, V. Chvátal (Eds.), Topics on Perfect Graphs, Annals of Discrete Mathematics, Vol. 21, North-Holland, Amsterdam, 1984, pp. 63–65) if there exists a linear order $<$ on its set of vertices such that no induced path $abcd$ with edges ab, bc, cd has both $a < b$ and $d < c$. Given a graph G and a vertex v in G such that $G - v$ is perfectly orderable, we set some conditions on v for which we deduce that G is perfectly orderable. Our method allows to construct a new class of such graphs, recognizable in polynomial time, containing quasi-brittle graphs, charming graphs and some other classes of perfectly orderable graphs. © 2001 Elsevier Science B.V. All rights reserved.

1. Introduction

A natural way to colour the vertices of a graph $G = (V, E)$ ordered in a sequence $v_1 < v_2 < \dots < v_n$ by the set of ‘colours’ $\{1, 2, 3, \dots\}$ consists of scanning the sequence from v_1 to v_n and assigning to each v_j the smallest integer assigned to none of its neighbours v_i such that $i < j$. This way of colouring the vertices of G is called *the greedy algorithm*. Certainly, this way may use a number of colours greater than the chromatic number of G . A graph G is said to be *perfectly orderable* [2] if there exists a linear order $<$ such that no induced P_4 $abcd$ in G has both $a < b$ and $d < c$ (such a P_4 is called an *obstruction* in $(G; <)$). Obviously, such an order, called *perfect order*, is also a perfect order on every $V' \subseteq V$, therefore the family of perfectly orderable graphs is hereditary. Recall that C. Berge defined a graph G to be *perfect* if the vertices of every induced subgraph H of G can be coloured with $\omega(H)$ colours, where $\omega(H)$ is the

* Corresponding author.

E-mail address: ht@lifo.univ-orleans.fr (H. Thuillier).




Fig. 1.

maximum clique size in H . It can be easily seen that a graph G is perfect if and only if every induced subgraph H contains a stable set intersecting every maximum clique in H . Berge and Duchet [1] defined a graph G to be *strongly perfect* if every induced subgraph H of G contains a stable set that intersects every maximal clique in H (as usual, ‘maximal’ is meant with respect to set-inclusion). Chvátal [2] has shown that perfectly orderable graphs are strongly perfect and that the greedy algorithm applied to a perfect order gives an optimal colouring.

For any integer $k \geq 3$, an induced path (respectively cycle) on k vertices will be denoted by P_k (respectively C_k). An induced subgraph of a graph G isomorphic to a P_k (respectively C_k) is simply said to be a P_k in G (resp. a C_k in G). A vertex v in a graph G is said to be *semi-simplicial* [6] if v is midpoint of no P_4 in G . A vertex v in a graph G is said to be *charming* [8] if v is not end-vertex in a P_5 in G , is not end-vertex in a P_5 in \bar{G} and does not lie on a C_5 in G .

A *rooted graph* is a pair $H=(F, u)$ where F denotes a graph and u a vertex in F . Let \mathcal{F} be a family of rooted graphs. A vertex v in a graph G is said to be \mathcal{F} -free if there is no induced subgraph F of G such that $H=(F, v)$ is isomorphic to a rooted graph of \mathcal{F} . For example, a semi-simplicial vertex is \mathcal{F} -free for the family $\mathcal{F}=\{(P_4, u)\}$ where u denotes a midpoint of P_4 , a charming vertex is \mathcal{F} -free for the family \mathcal{F} described in Fig. 1.

Hoàng et al. [8] call *charming* any graph in which every induced subgraph has a charming vertex, and prove that every charming graph is perfectly orderable. A natural idea is to consider the following related notion: a vertex v in a graph G is said to be *nice* if v is not internal vertex in a P_5 in G , is not internal vertex in a P_5 in \bar{G} and does not lie on a C_5 in G . A nice vertex is \mathcal{F} -free for the family \mathcal{F} described in Fig. 2. We call *nice* any graph in which every induced subgraph has a nice vertex, and we will prove that every nice graph is perfectly orderable (see below). In fact, this result is a corollary of Theorem 3 in Section 5.

An edge ab in a graph G is called a *symmetric wing* if there exist vertices c, d, p, q such that both $abcd$ and $baqp$ are induced P_4 ’s in G . A vertex v in G is said to be *special* [10] if v is incident with no symmetric wing in G and \bar{G} .

Lemma 1. *An edge ab in a graph G is a symmetric wing if and only if there is an induced subgraph of G depicted in Fig. 3 such that ab is the edge whose ends are circled.*




Fig. 2.




Fig. 3.

Proof. Let c, d, p and q such that both $abcd$ and $baqp$ are P_4 's in G . There are two cases according as $d = p$ or $d \neq p$. If $d = p$ then cq is the only one possible edge. If $d \neq p$ then cq, cp, dq, dp are possible edges. By examining every case, it is a routine matter to obtain the announced result. \square

Lemma 2. A vertex v in a graph G is special if and only if v is \mathcal{F}_1 -free, where \mathcal{F}_1 is the family of rooted graphs $\{H_1, H_2, \bar{H}_2, H_4, \bar{H}_4\}$ (see Figs. 2 and 4).

Proof. By considering an end-vertex a of a symmetric wing and the rooted graphs (C_5, a) , (\bar{P}_5, a) and (A, a) in Fig. 3 and their (rooted) complements, we obtain rooted graphs isomorphic to H_1, H_2, \bar{H}_2, H_4 or to \bar{H}_4 (see Figs. 2 and 4). The other graphs in Fig. 3, that is (P_6, a) , (C_6, a) , (B, a) , and their complements give rooted graphs containing a rooted graph (F, a) isomorphic either to H_2 or to \bar{H}_2 depicted in Fig. 2.

Conversely, let v be a vertex belonging to a rooted graph H of \mathcal{F}_1 . If H belongs to $\{H_1, \bar{H}_2, H_4\}$ (resp. $\{H_2, \bar{H}_4\}$) then v is end-vertex in a symmetric wing in G (resp. \bar{G}). \square




Fig. 4.

In [10] Olariu defined a graph G to be *quasi-brittle* if every induced subgraph H of G contains a special vertex, and proved that such a graph is perfectly orderable. Indeed, an attentive examination of his proof shows that he obtained the following more general result:

Theorem 1. *Let G be a graph and v be a special vertex in G . Then, G is perfectly orderable if and only if $G - v$ is perfectly orderable.*

Note that such a result was known if v is semi-simplicial [6] or if v is a charming vertex [8].

Our purpose is to generalize these previous results and to obtain new families of perfectly orderable graphs by proving statements of the following type:

Assertion A. *Let \mathcal{F} be a family of rooted graphs. Let G be a graph and v be a \mathcal{F} -free vertex in G . Then, G is perfectly orderable if and only if $G - v$ is perfectly orderable.*

2. Definitions and notations

For terms not defined here, the reader is referred to [5]. All the graphs in this paper are simple and a graph is denoted $G = (V(G), E(G))$ (or simply $G = (V, E)$ if no confusion exists); its *complement* \bar{G} is the graph (V, \bar{E}) where $\bar{E} = \{xy \mid x \in V, y \in V \text{ and } xy \notin E\}$. For any subset A of V , the subgraph of G (resp. \bar{G}) induced by the set A will be denoted by $G[A]$ (resp. $\bar{G}[A]$).

For any vertex v in V , the *neighbourhood* $N_G(v)$ of v in G is defined as the set of all vertices adjacent to v in G . These vertices are called *neighbours* of v . When there is no possibility of confusion, the notation $N(v)$ will also be used to denote the neighbourhood of v . The set of vertices in G which are not adjacent to v is denoted by $M(v)$ (this set is equal to $N_{\bar{G}}(v)$, the neighbourhood of v in \bar{G}). For any subset W of $V \setminus \{v\}$ such that $W \cap N(v) \neq \emptyset$ and v is not adjacent to at least one vertex in W , we will say that v *sees* W *partially*. If $W \subseteq N(v)$, we will say that v *sees* W *completely*.

If $A \subseteq V$, then $G \setminus A$ will denote the graph induced by $V \setminus A$. The particular case where $A = \{x\}$ will be denoted $G - x$; also if H is a subgraph of G , we denote by $H + x$ the subgraph induced in G by $V(H) \cup \{x\}$ (if $x \notin V(H)$).

For any path P , the *length* of P is the number of its edges. If $V(P) = \{v_1, \dots, v_k\}$ and $E(P) = \{v_i v_{i+1} \mid i \in \{1, \dots, k-1\}\}$, P is also denoted by $[v_1, \dots, v_k]$. The vertices v_1 and v_k are its *end-vertices* while any vertex v_i , with $1 < i < k$, is said to be an *internal* vertex. An induced path on k vertices will be denoted by P_k . For a P_4 with vertices a, b, c, d (in this order on the path), the notation will be slightly changed: it will be denoted $abcd$. The two internal vertices b and c will be called *midpoints* while the end-vertices a and d will be also referred as *endpoints*.

Similarly, a chordless cycle on k vertices is denoted by C_k or by $[v_1, \dots, v_k, v_1]$ if its vertex set is $\{v_1, \dots, v_k\}$ and its edge set is $\{v_i v_{i+1} \mid 1 \leq i \leq k-1\} \cup \{v_k v_1\}$.

A set $A \subseteq V$ of vertices is called a *module* if every vertex in $V \setminus A$ is either adjacent to all vertices in A , or none of them. A module in G is also a module in \bar{G} . A module is an *homogeneous set* if it is not a *trivial* one, i.e. not an empty set, a singleton or V itself. A graph with more than two vertices is *prime* if it only has trivial modules. Note that P_4 is the smallest prime graph and that any prime graph distinct from P_4 has at least five vertices.

Let G and H be two graphs and let v be a vertex in G . The χ -join of H and G into v is the graph obtained from the union of H and $G - v$ by adding all the edges xy with $x \in V(H)$ and $y \in N_G(v)$; this graph is denoted by $\chi(H, G; v)$. Since an induced P_4 in $\chi(H, G; v)$ not contained in H and not in $G - v$ has exactly one vertex in H and the three other vertices in $G - v$, note that if $v_1 < v_2 < \dots < v_n$ is a perfect order of G and if $w_1 < w_2 < \dots < w_p$ is a perfect order of H , then $v_1 < \dots < v_{i-1} < w_1 < w_2 < \dots < w_p < v_{i+1} < \dots < v_n$ is a perfect order of $\chi(H, G; v_i)$. Then we have

Lemma 3 (Chvátal [3]). *Let G and H be two perfectly orderable graphs and let v be a vertex in G . Then $\chi(H, G; v)$ is perfectly orderable.*

3. Pleasant vertices

A vertex v in a graph G is said to be a *pleasant vertex* if v is \mathcal{F}_0 -free, with $\mathcal{F}_0 = \{H_1, H_2, \bar{H}_2\}$ where these rooted graphs are described in Fig. 2. Note that special vertices and nice vertices are pleasant vertices. We call *pleasant* any graph in which every induced subgraph has a pleasant vertex.

For any vertex v in a graph G let $S_1(v), \dots, S_{q_v}(v)$ be the connected components of $\bar{G}[N(v)]$, and let $Q_1(v), \dots, Q_{k_v}(v)$ be the connected components of $G[M(v)]$. For the sake of simplicity we will denote these components by S_1, \dots, S_q and by Q_1, \dots, Q_k .

We give here a structural result that we will often use.

Proposition 1. *Let v be a pleasant vertex in a graph $G=(V, E)$. For every $i \in \{1, \dots, q\}$ and every $j \in \{1, \dots, k\}$ one of the following statement is true:*

- (1) every vertex in S_i is adjacent to every vertex in Q_j ,
- (2) every vertex in S_i has no neighbour in Q_j ,

- (3) there exists a non trivial part A_i of S_i such that for every vertex x in S_i and for every vertex y in Q_j , xy belongs to E if and only if x belongs to A_i ,
 (4) there exists a non trivial part B_j of Q_j such that for every vertex x in S_i and for every vertex y in Q_j , xy belongs to E if and only if y belongs to B_j .

Proof. If $|S_i| = 1$ or $|Q_j| = 1$ then the result is clearly true. We suppose that $|S_i| \geq 2$, $|Q_j| \geq 2$ and that none of the conditions (1)–(3) is satisfied. Then, there exist $x \in S_i$, a and $b \in Q_j$ such that $xa \in E$ and $xb \notin E$. Since Q_j is connected in G , we can suppose without loss of generality that $ab \in E$. Since S_i is connected in \bar{G} , there is $y \in S_i$ distinct from x such that $xy \notin E$. Since the set of vertices $\{y, v, x, a, b\}$ induces no cycle C_5 , no P_5 and no \bar{P}_5 , we have $ya \in E$ and $yb \notin E$.

Now we want to prove that any vertex h in Q_j is either adjacent to every vertex in S_i or adjacent to no vertex in S_i . A contrario, suppose that there exist h in Q_j , c and d in S_i such that $hc \in E$ and $hd \notin E$. Since S_i is connected in \bar{G} , we may suppose without loss of generality that $cd \notin E$. Let k be a neighbour of h in Q_j . Since the set $\{k, h, c, v, d\}$ induces no C_5 , no P_5 and no \bar{P}_5 , we have $kc \in E$ and $kd \notin E$. Since Q_j is connected, c is adjacent to every vertex in Q_j (and d is adjacent to no vertex in Q_j). But b belongs to Q_j and is adjacent to no vertex in S_i , so we obtain a contradiction. Thus, the statement 4) is true. \square

Corollary 1. Let v be a pleasant vertex in a graph $G = (V, E)$.

- (i) Let S_i be a component of $\bar{G}[N(v)]$. Let a and b be two vertices in $M(v)$ such that $N(a) \cap S_i \neq \emptyset$, $N(b) \cap S_i \neq \emptyset$ and $N(a) \cap S_i \neq N(b) \cap S_i$. Then a and b belong to distinct components of $G[M(v)]$.
 (ii) Let Q_j be a component of $G[M(v)]$. Let a and b be two vertices in $N(v)$ such that $N(a) \cap Q_j \neq \emptyset$, $N(b) \cap Q_j \neq \emptyset$ and $N(a) \cap Q_j \neq N(b) \cap Q_j$. Then a and b belong to distinct components of $\bar{G}[N(v)]$.

4. Acyclic orientation and obstructions

A perfect order of G can be seen as an acyclic orientation of the edges of G such that every induced P_4 , $abcd$, is not an obstruction, that is $abcd$ cannot have simultaneously (a, b) and (d, c) as arcs. Our aim is the following: given a perfect order $<$ of $G - v$, how to orient the edges incident with v and how to modify the orientation of some edges in $G - v$ so that the oriented graph G has no circuit and no obstruction. We may define a linear order $<'$ of G in the following way: let us partition $V - v$ into L and R , and denote $N(v) \cap L$ by LH , $M(v) \cap L$ by LL , $N(v) \cap R$ by RH and $M(v) \cap R$ by RL ; we do not modify the orientations in $G[L]$ and $G[R]$, every edge xy such that $x \in L$ and $y \in R$ is oriented such that $x <' y$, every edge xv with $x \in L$ is oriented such that $x <' v$ and, at last, every edge yv such that $y \in R$ is oriented such that $v <' y$ (see Fig. 5). Now, our problem is to define the sets LH , LL , RH and RL in a suitable way to avoid obstructions.




Fig. 5.

We note that, to obtain their result, Hoàng et al. [8] set $LH = N(v)$, $LL = \emptyset$, $RH = \emptyset$ and $RL = M(v)$.

We consider a pleasant vertex v of the graph G , and we suppose $G - v$ perfectly orderable. Let $<$ be a perfect order on $G - v$. Following a similar idea as used by Olariu [10], a component S_i of $\tilde{G}[N(v)]$ is called *impure* if a vertex a in $M(v)$ and vertices b, d in S_i exist such that $abvd$ is a P_4 and $a < b$. A component S_i is called *pseudo-pure* if there are vertices a, b in $M(v)$ and a vertex c in S_i such that $abcv$ is a P_4 and $a < b$. A component which is neither impure nor pseudo-pure is referred to as *pure*.

Lemma 4. *No pseudo-pure component is impure.*

Proof. Assume that S_i is both pseudo-pure and impure: let a, b, x in $M(v)$ and c, y, z in S_i be such that $abcv$ and $xyvz$ are P_4 with $a < b$ and $x < y$. Since $ac \notin E$ and $bc \in E$, the vertex b sees S_i completely (by Proposition 1). As $bz \in E$, $xy \in E$ and $xz \notin E$, the component Q_j containing x is different from the component containing a and b (by Corollary 1). Then $abyx$ is an obstruction in $(G - v; <)$, a contradiction. \square

We can now define the sets LH , RH , LL and RL : LH contains all the pseudo-pure components and, perhaps, some pure components; RH contains all the impure components and the remaining pure components; LL is the union of some components Q_j and RL is the union of the other components of $G[M(v)]$ (LL or RL can be empty). By construction, the oriented graph $(G; <')$ presents no circuit.

Claim 1. $(G; <')$ presents no obstruction containing v .

Proof. If $abcv$ is an obstruction in $(G; <')$, then a and b belong to the same component Q_j and $a < b$. Thus, c belongs to a pseudo-pure component and $v <' c$, a contradiction. If $abvd$ is an obstruction in $(G; <')$, then the component S_i containing b and d is




Fig. 6.

contained in LH ($d <' v$) and a is in LL ($a <' b$). Thus $a < b$ and S_i is impure, a contradiction. \square

We are going to observe now the possible obstructions. Let $abcd$ be an obstruction in $(G; <')$. Without loss of generality, we can suppose that $a \in L$, $b \in R$ and $b < a$. Since there is no edge between LL and RL , we notice that a or b is a neighbour of the vertex v .

Proposition 2. *If $abcd$ is an obstruction in $(G; <')$, then one of the following situations depicted in Fig. 6 is realized:*

Situation 1: $a \in LH$ and $b, c, d \in Q_j \subseteq RL$.

Situation 2: $a \in LL$, $b, c \in RH$ and $d \in RL$.

Situation 3: $a \in LL$ and $b, c, d \in RH$.

Situation 4: $a \in LL$, $b, d \in RH$ and $c \in RL$.

Situation 5: $a, d \in LL$ and $c, b \in RH$.

Situation 6: $a, d \in LL$, $c \in LH$ and $b \in RH$.

Proof. Since $d <' c$, we notice that $c \notin L$ while $d \in R$. We distinguish three cases:

Case A: $c, d \in R$;

Case B: $d \in L$ and $c \in R$;

Case C: $c, d \in L$.

Case A: $c, d \in R$, then $d < c$.

Case A1: We suppose that $a, b \in N(v)$. Let S_i be the component (pure or pseudo-pure) containing a and let S_j be the component (pure or impure) containing b . Since $ad \notin E$ and $bd \notin E$, d belongs to $M(v)$. If $c \in N(v)$ then c belongs to S_i and this component is impure (because $d < c$), a contradiction. Thus, c and d belong to $M(v)$. But now, we consider the P_4 $dcbv$ and we see that S_j is pseudo-pure, which gives also a contradiction. Then, Case A1 does not occur.

Case A2: We suppose that $a \in N(v)$ and $b \notin N(v)$. As $ac, ad \notin E, b, c$ and d belong to the same component Q_j . This is the first situation.

Case A3: We suppose that $a \notin N(v)$ and $b \in N(v)$. As the component containing b is not pseudo-pure, one of the vertices c or d is in $N(v)$. We obtain three other possibilities: Situations 2–4.

Case B: $d \in L$ and $c \in R$.

Case B1: We suppose that $a \in N(v)$. Since $ac, bd \notin E$ and $cd \in E$, c belongs to $M(v)$ and d belongs to $N(v)$. Since $ad, bd \notin E$, a and d are in the same component S_i , b and c are in the same component Q_j . By Proposition 1, $abcd$ cannot be a P_4 ; this is a contradiction. Then, Case B1 cannot occur.

Case B2: We suppose that $a \notin N(v)$ and $b \in N(v)$. Since $bd \notin E$ and $cd \in E$, we have $d \notin N(v)$ and $c \in N(v)$. We obtain the fifth situation.

Case C: $c, d \in L$, then $d < c$.

Case C1: We suppose that $a, b \in N(v)$. As $bd \notin E$, necessarily $d \notin N(v)$. If $c \in N(v)$, then c belongs to the component S_i containing a , and S_i is impure; this is a contradiction. If $c \notin N(v)$, we consider the graph induced by $\{d, c, b\}$ and we see that the component $S_{i'}$ containing b is pseudo-pure. This is also a contradiction; then Case C1 is impossible.

Case C2: We suppose that $a \in N(v)$ and $b \notin N(v)$. Necessarily, a and c are in the same component S_i . If $d \in N(v)$, then d is in the component S_i and $bavd$ is a P_4 where $b < a$. Then S_i is impure, which is impossible. If $d \notin N(v)$, then $dcva$ is a P_4 and $d < c$. So the component S_i is impure. Thus, Case C2 is impossible.

Case C3: We suppose that $a \notin N(v)$ and $b \in N(v)$. If $c \notin N(v)$, then $dcbv$ is a P_4 , where $d < c$, and the component containing b is pseudo-pure. So $c \in N(v)$, and we obtain the last situation. \square

From the previous proposition, we deduce immediately the following two results (we shall see later that we can improve Corollary 3).

Corollary 2. *Let v be a pleasant vertex of a graph G . If v is endpoint of no P_5 then G is perfectly orderable if and only if $G - v$ is perfectly orderable.*

Proof. We choose $LL = \emptyset$ and $RL = M(v)$: none of Situations 2–6 appears. Since v is endpoint of no P_5 , Situation 1 cannot be realized. \square

Corollary 3. *Let v be a pleasant vertex of a graph G . If v is $\{H_5, H_6\}$ -free (H_5 and H_6 are described in Fig. 7), then G is perfectly orderable if and only if $G - v$ is perfectly orderable.*




Fig. 7.

Proof. We choose $LL = M(v)$ and $RL = \emptyset$; the Situations 1, 2, 4 cannot appear. Since v is a vertex $\{H_5, H_6\}$ -free, none of the Situations 3, 5, and 6 is realized. \square

5. Main results

Let us consider again the following

Assertion A. Let \mathcal{F} be a family of rooted graphs. Let G be a graph and v be a \mathcal{F} -free vertex in G . Then, G is perfectly orderable if and only if $G - v$ is perfectly orderable.

We have solely to prove that if $G - v$ is perfectly orderable then G is perfectly orderable. As a consequence of Lemma 3, we have:

Claim 2. To prove Assertion A by induction on the number of vertices, it can be supposed that G is prime.

Proof. If every graph in \mathcal{F} has p vertices or more, then this result is true for any graph G such that $|V(G)| < \min(4, p)$. Let us suppose that for any graph H such that $|V(H)| < |V(G)|$, H is perfectly orderable if w is a \mathcal{F} -free vertex in H and $H - w$ is perfectly orderable. We suppose that G contains a homogeneous set M .

(i) If $v \notin M$, Let H be the graph obtained from G by contracting M in a single vertex z . Since H is isomorphic to an induced subgraph of G , v is a \mathcal{F} -free vertex in H . Since $H - v$ is an induced subgraph of $G - v$, $H - v$ is perfectly orderable. By induction, H is perfectly orderable. The subgraph $G[M]$ is an induced subgraph of $G - v$, thus it is a perfectly orderable graph. By Lemma 3, $G = \chi(G[M], H; z)$ is perfectly orderable.

(ii) If $v \in M$ then $G[M] - v$ is an induced subgraph of $G - v$, thus $G[M] - v$ is perfectly orderable. Since v is a \mathcal{F} -free vertex in $G[M]$, by induction $G[M]$ is perfectly orderable. The graph $H = G - M + v$ is isomorphic to $G - M + u$ with $u \in M - v$, a subgraph of $G - v$, thus H is perfectly orderable. By Lemma 3, $G = \chi(G[M], H; v)$ is perfectly orderable. \square

The following statement generalizes Theorem 1.

Theorem 2. *Let G be a graph and v be a vertex \mathcal{F}_2 -free, where \mathcal{F}_2 is the family $\{H_1, H_2, \tilde{H}_2, \tilde{H}_4\}$ (see Figs. 2 and 4). Then, G is perfectly orderable if and only if $G - v$ is perfectly orderable.*

Lemma 5. *Let G be a prime graph and v be a \mathcal{F}_2 -free vertex in G . If S_i is a component of $\tilde{G}[N(v)]$ and u is a non-isolated vertex in $G[M(v)]$, then u is adjacent to every vertex of S_i or none.*

Proof. On the contrary, we suppose that the vertex u sees partially the component S_i : there are vertices a, b in S_i such that $au \in E$ and $bu \notin E$. Without loss of generality, we can suppose that $ab \notin E$. As G is prime, the component Q_j containing u is not an homogeneous set, and there are vertices x in $N(v)$ and y, z in Q_j such that $xy \in E$ and $xz \notin E$. We can suppose that $yz \in E$. By Proposition 1, we have $ay \in E$, $az \in E$, $by \notin E$, $bz \notin E$, and by Corollary 1, $x \notin S_i$. Then, the rooted graph $(G[v, a, b, y, z, x], v)$ is isomorphic to \tilde{H}_4 . \square

Lemma 6. *Let v be a pleasant vertex in a graph G , such that $G - v$ is perfectly orderable. Let $<$ be a perfect order of $G - v$, S_i be an impure component and x be a vertex in $M(v)$ which sees S_i completely. Then there exists a vertex $y \in S_i$ such that $x < y$.*

Proof. Let S_i be an impure component. There exist vertices $a \in M(v)$ and $b, y \in S_i$ such that $abvy$ is a P_4 and $a < b$. If $x \in M(v)$ sees S_i completely then, by Corollary 1, x and a are not in the same component of $G[M(v)]$. Since $abxy$ is not an obstruction in $(G - v; <)$, we have $x < y$. \square

Proof of Theorem 2. By Claim 2, we can suppose that G is prime. We choose LH as the union of all the pure or pseudo-pure components, RH as the union of all the impure components, RL as the set of all the isolated vertices in $G[M(v)]$ and LL as the union of the other components of $G[M(v)]$.

Claim 3. *None of the Situations 1, 3, 4 appears.*

Proof. In Situation 1, vertices b, c, d are not isolated in $G[RL]$. In Situations 3 and 4, the vertex a belongs to $M(v)$ and sees partially the impure component containing the vertices b and d . By Lemma 5, a is isolated in $G[M(v)]$ and cannot belong to LL . \square

Claim 4. *Situations 5 and 6 are not realized.*

Proof. In Situations 5 and 6, a and d are not isolated vertices in $G[M(v)]$. Let S_i (resp. S_j) be the impure component containing b (resp. c). By Lemma 5, a (resp. d)

sees completely S_i (resp. S_j). Since $ac \notin E$, S_i and S_j are distinct components. By Lemma 6, in Situation 5, there exist vertices $c', b' \in RH$ such that $dc'b'a$ is an obstruction in $(G - v; <)$; in Situation 6, there exists a vertex $b' \in RH$ such that $dc'b'a$ is an obstruction in $(G - v; <)$. \square

A careful reading of Olariu's study about quasi-brittle graphs [10, Proof of Theorem 2, pp. 151–152] allows us to write the following lemma:

Lemma 7 (Olariu [10]). *Let v be a \mathcal{F}_2 -free vertex in a graph G , such that $G - v$ is perfectly orderable. There exists no obstruction $abcd$ in $(G; <')$, where $a \in LL$ and $d \in RL$.*

Thus, we have

Claim 5. *Situation 2 does not appear.*

This ends the proof of Theorem 2. \square

Another way to generalize Theorem 1 is to consider the family $\{H_1, H_2, \bar{H}_2, H_4\}$. Unfortunately, we are unable to prove the corresponding result (we must add the rooted graph H_8 defined below; see Corollary 6).

Theorem 3. *Let G be a graph and v be a \mathcal{F}_3 -free vertex, where \mathcal{F}_3 is the family of rooted graphs $\{H_1, H_2, \bar{H}_2, H_7, H_8\}$ (see Figs. 2 and 8). Then, G is perfectly orderable if and only if $G - v$ is perfectly orderable.*

Proof. Let $<$ be a perfect order on $G - v$. Since v is a pleasant vertex of G , by Lemma 4, no pseudo-pure component is impure. Then, let us consider the linear order $<'$ on G defined in Section 4, such that LH contains no pure component (that is LH is the union of the pseudo-pure components) and LL is empty (that is $RL = M(v)$). The oriented graph $(G; <')$ presents no circuit and, by Claim 1, no obstruction containing v . Let $A = abcd$ be an obstruction in $(G; <')$. Since $LL = \emptyset$, by Proposition 2, Situation 1 is the only one possibility. We suppose without loss of generality that a belongs to




Fig. 8.

LH and b, c and d belong to RL . We denote by $C(A)$ the pseudo-pure component of $\bar{G}[N(v)]$ containing a and by $Q(A)$ the component of $G[M(v)]$ containing b, c and d . Note that, by Proposition 1, for every $x \in C(A)$ we have $bx \in E$, $cx \notin E$ and $dx \notin E$. Moreover, $a <' b$, $b < a$ and $d < c$.

Claim 6. *There exist s and t in $Q(A)$ such that $t < s$ and for all x in $C(A)$, $sx \in E$ and $tx \notin E$.*

Proof. Since $C(A) \subseteq LH$, $C(A)$ is a pseudo-pure component. Then, there exist t and s in a component Q_j such that $t < s$, $st \in E$ and for all x in $C(A)$, $sx \in E$ and $tx \notin E$. If $Q_j \neq Q(A)$ then $tsab$ is an obstruction in $(G - v; <)$, a contradiction. Thus, $Q_j = Q(A)$. \square

Let \mathcal{A} be the set of obstructions in $(G; <')$ and let $\mathcal{C}(\mathcal{A})$ be the set of vertices x in R such that there exist $A = abcd$ in \mathcal{A} and an oriented path, $P[x, b]$, from x to b , in $(G - v; <)$. Note that there is no edge xy such that $x \in \mathcal{C}(\mathcal{A})$, $y \in R \setminus \mathcal{C}(\mathcal{A})$ and $y < x$.

Now, set $L' = L \cup \mathcal{C}(\mathcal{A})$ and $R' = R \setminus \mathcal{C}(\mathcal{A})$. We apply to this new partition $\{L', R'\}$ of $V \setminus \{v\}$ the rules given in Section 3 to obtain from $(G - v; <)$ a new oriented graph $(G; <'')$.

Remark 1. If there exists xy in E such that $y <' x$ and $y > x$, then $x \in R'$ and $y \in LH = L'H \setminus \mathcal{C}(\mathcal{A})$.

Note that, by Remark 1, Situations 2–6 do not appear in $(G - v; <'')$.

Remark 2. By definition of $\mathcal{C}(\mathcal{A})$, Situation 1 does not appear in $(G - v; <'')$.

Clearly, $(G; <'')$ presents no circuit. Remarks 1 and 2 imply that, if B is an obstruction in $(G; <'')$ then B contains v .

If $B = vv_2v_3v_4$ then $v_2 \in R'H$ (otherwise $v_2 <' v$). Clearly, v_3 and v_4 belong to $M(v)$ and by Remark 1, $v_4 < v_3$. But the component S_i containing v_2 is pseudo-pure and then $v_2 \in LH \subseteq L'H$, a contradiction.

If $B = v_1vv_2v_3$ then $v_1 \in L'H$ (otherwise $v <' v_1$). Clearly, v_1 and v_2 are in the same component S_i and $v_3 \in M(v)$. Since $v_3 <' v_2$, by Remark 1, $v_3 < v_2$. Hence, S_i is an impure component, that is $v_1 \notin LH$. Since $v_1 \in L'H$, $v_1 \in \mathcal{C}(\mathcal{A})$.

Thus, there exist $A = abcd$ in \mathcal{A} and an oriented path in $(G - v; <)$, $P[v_1, b]$, from v_1 to b . Since $C(A) \subseteq LH$, $C(A) \neq S_i$, $av_1 \in E$ and $av_2 \in E$. Since $b < a$, $v_1 < a$ (otherwise $v_1P[v_1, b]bav_1$ is a circuit in $(G - v; <)$). Moreover, $v_3a \in E$, otherwise $v_1av_2v_3$ would be an obstruction in $(G - v; <)$.

We distinguish two cases:

Case A: $v_1b \in E$. Either $P[v_1, b] = v_1b$ or $v_1P[v_1, b]bv_1$ is a cycle. In each case $v_1 < b$. By Corollary 1, $v_3 \notin Q(A)$. Then $v_3b \notin E$. Note that $v_2b \notin E$, otherwise $v_1bv_2v_3$ would

be an obstruction in $(G - v; <)$. By Proposition 1, v_1 is adjacent to every vertex in $Q(A)$ and v_2 is adjacent to no vertex of $Q(A)$. Now, the subgraph of G induced by $\{v, a, v_1, v_2, v_3, b, c, d\}$ is isomorphic to the rooted graph H_8 depicted in Fig. 8, a contradiction.

Case B: $v_1b \notin E$. By Claim 6 there exist s and t in $Q(A)$ such that $sa \in E$, $ta \notin E$ and $t < s$. Since v_1ast is not an obstruction in $(G - v; <)$, $v_1s \in E$ or $v_1t \in E$. Then, by Proposition 1, v_1 and v_2 have the same neighbourhood in $Q(A)$. Thus, by Corollary 1, $v_3 \notin Q(A)$.

Case B1: If $v_1t \in E$. Clearly $t \neq b$ and $v_2t \in E$. Since $v_1tv_2v_3$ cannot be an obstruction, we have necessarily $t < v_1$. Moreover $tb \in E$, otherwise tv_1ab would be an obstruction in $(G - v; <)$. But now, the subgraph of G induced by $\{v, a, v_1, v_2, v_3, b, t\}$ is isomorphic to the rooted graph H_7 depicted in Fig. 8, a contradiction.

Case B2: If $v_1t \notin E$. Then necessarily, $v_1s \in E$ and $v_2s \in E$. Since tsv_2v_3 is not an obstruction in $(G - v; <)$, $v_2t \in E$. Then $v_1t \in E$, a contradiction. \square

From Theorem 3, we deduce the following three corollaries:

Corollary 4. *Let G be a graph and v be a nice vertex in G (see Fig. 2). Then, G is perfectly orderable if and only if $G - v$ is perfectly orderable.*

Proof. The rooted graph H_3 is a rooted subgraph of H_8 and \bar{H}_3 is a rooted subgraph of H_7 . \square

Corollary 5. *Let G be a graph and v be a \mathcal{F}_4 -free vertex, where \mathcal{F}_4 is the family of rooted graphs $\{H_1, H_2, \bar{H}_2, H_5\}$ (see Figs. 4 and 7). Then, G is perfectly orderable if and only if $G - v$ is perfectly orderable.*

Proof. The rooted graph H_5 is a rooted subgraph of H_7 and H_8 . \square

Corollary 6. *Let G be a graph and v be a \mathcal{F}_5 -free vertex, where \mathcal{F}_5 is the family of rooted graphs $\{H_1, H_2, \bar{H}_2, H_4, H_8\}$ (see Figs. 2, 4 and 7). Then, G is perfectly orderable if and only if $G - v$ is perfectly orderable.*

Proof. The rooted graph H_4 is a rooted subgraph of H_7 . \square

Remark 3. Since H_4 is a rooted subgraph of H_7 and \bar{H}_4 is a rooted subgraph of H_8 , we note that Theorem 1 is also a corollary of Theorem 3.

6. Conclusion

Chvátal [3] defined a graph G to be *brittle* if each induced subgraph F of G contains a vertex that is not a midpoint of any P_4 (semi-simplicial vertex) or not an endpoint of any P_4 . If v is not a midpoint of any P_4 or not an endpoint of any P_4 and if $F - v$ is

perfectly orderable then F is perfectly orderable. Thus, every brittle graph is perfectly orderable [3] (see also [7]).

Let $\{\mathcal{G}_i\}_{1 \leq i \leq k}$ be a finite set of finite families of rooted graphs such that for any $i \in \{1, \dots, k\}$ Assertion A is true. For example, $\mathcal{G}_1 = \{(P_4, v)\}$ where v denotes a midpoint of P_4 , $\mathcal{G}_2 = \{(P_4, v)\}$ where v denotes an endpoint of P_4 , \mathcal{G}_3 is the family depicted in Fig. 1 (\mathcal{G}_3 -free vertices are charming vertices), \mathcal{G}_4 is the family in Fig. 2 (\mathcal{G}_4 -free vertices are nice vertices), $\mathcal{G}_5 = \mathcal{F}_2$, $\mathcal{G}_6 = \mathcal{F}_3$. We define a class $\mathcal{P}(\mathcal{G}_1, \dots, \mathcal{G}_k)$ of perfectly orderable graphs in the following way: a graph G belongs to $\mathcal{P}(\mathcal{G}_1, \dots, \mathcal{G}_k)$ if for any subgraph F of G there exist a family \mathcal{G}_i and a \mathcal{G}_i -free vertex v in F . According to the previous example, we note that $\mathcal{P}(\mathcal{G}_1, \mathcal{G}_2)$ is the class of brittle graphs and that $\mathcal{P}(\mathcal{G}_1, \dots, \mathcal{G}_6) = \mathcal{P}(\mathcal{G}_3, \dots, \mathcal{G}_6)$. The last class contains brittle graphs, charming graphs, nice graphs and quasi-brittle graphs.

Although recognizing general perfectly orderable graphs is NP-complete [9], clearly, there is a polynomial time recognition algorithm for $\mathcal{P}(\mathcal{G}_1, \dots, \mathcal{G}_k)$.

One may ask the following question: is it true that if v is a pleasant vertex in a graph G such that $G - v$ is perfectly orderable, then G is perfectly orderable? In [4] we show a graph giving a negative answer to this question.

References

- [1] C. Berge, P. Duchet, Strongly Perfect Graphs, in: C. Berge, V. Chvátal (Eds.), Topics on Perfect Graphs, Annals of Discrete Mathematics, Vol. 21, North-Holland, Amsterdam, 1984, pp. 57–61.
- [2] V. Chvátal, Perfectly ordered graphs, in: C. Berge, V. Chvátal (Eds.), Topics on Perfect Graphs, Annals of Discrete Mathematics, Vol. 21, North-Holland, Amsterdam, 1984, pp. 63–65.
- [3] V. Chvátal, Perfect Graph Seminar, McGill University, Montreal, Canada, 1983.
- [4] J.L. Fouquet, F. Roussel, P. Rubio, H. Thuillier, On pleasant vertices in graphs, Report RR-99-14, LIFO, University of Orleans, 1999.
- [5] M.C. Golumbic, Algorithmic Graph Theory and Perfect Graphs, Academic Press, New York, 1980.
- [6] B. Jamison, S. Olariu, On the semi-perfect elimination, Adv. Appl. Math. 9 (1988) 364–376.
- [7] C.T. Hoàng, N. Khouzam, On brittle graphs, J. Graph Theory 12 (3) (1988) 391–404.
- [8] C.T. Hoàng, F. Maffray, S. Olariu, M. Preissmann, A charming class of perfectly orderable graphs, Discrete Math. 102 (1992) 67–74.
- [9] M. Middendorf, F. Pfeiffer, On the complexity of recognizing perfectly orderable graphs, Discrete Math. 80 (1990) 327–333.
- [10] S. Olariu, Quasi-brittle graphs, a new class of perfectly orderable graphs, Discrete Math. 113 (1992) 143–153.